



# INSECTIDUST


Reg. No. / Nr.: Act / Wet 36 /1947

**A Dusting Powder Insecticide for use in Agriculture and Public Health with stomach and contact poison action for the control of pests as listed on Cotton, Groundnuts, Sorghum, Maize, Ornamentals, Sub-tropical fruits, Potatoes, Deciduous Fruit, Grazing, Wheat, Lucerne, Cruciferae, Mushrooms, Onions, Tobacco, Berans. Also for the control of Flies, Fleas and Bedbugs.**

**'n Stuifpoeier Insekmiddel vir gebruik in die Landbou en Openbare Gesondheid met maag- en kontakgif vir die beheer van plaes soos aangedui op Katoen, Grondbootjies, Graansorghum, Mielies, Sierstruik, Sub-tropiese vrugte, Aartappels, Sagte vrugte, Weidings, Koring, Lusern, Koolgewasse, Sampioene, Uie, Tabak, Boontjies. Ook vir die beheer van Vlieë, Vlooië en Weeluisse.**

Active ingredient : Mercaptothion (Organophosphate) .. 50 g/kg ...Merkaptotien (Organofosfaat)  
Pyrethrins ... 0,4 g/kg...Piretriene : Aktiewe bestanddeel

## Contents :



Registered by: AVIMA (PTY) LTD : Geregistreer deur  
P.O. BOX / POSBUS 3131  
KENMARE 1745  
Co. Reg. No. / Mpy Reg. Nr.: 1961 /01744 /07

Manufactured & formulated by: AVIMA (PTY) LTD :Vervaardig & geformuleer deur  
18 Aschenberg Street  
Chamdor, Krugersdorp

Batch No: : Lotnommer  
Date of Manufacture: :Vervaardigings datum

UN No: 2783 :UN Nr.  
EMERGENCY NUMBER: 0800 172 743 :NOODNOMMER



Caution  
Versigtig

**WARNINGS:**

- Do not dust within the following number of days, before harvest:

Curcurbits, Tomatoes and mushrooms = 1

Other edible crops = 7

- Do not apply to cucurbits unless plants are dry.
- Do not use where milk is processed or stored.
- Poisonous by contact, swallowing and inhalation.
- Toxic to fish, bees and wild life.
- Store away from food and feedstuffs.
- Keep out of reach of children, uninformed persons and animals.
- Keep container well closed.

ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL CLIMATIC AND STORAGE CONDITIONS; COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURRENCE OF RESISTANCE OF THE PEST AGAINST THE REMEDY CONCERNED AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHERMORE DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

**PRECAUTIONS:**

- Do not inhale dust.
- Wash with soap and water after use.
- Do not eat, drink or smoke whilst applying, or before washing hands and face or change of clothing.
- Prevent drift onto other crops, grazing, rivers, dams or areas not under treatment.
- Prevent contamination of food, feedstuffs and water.
- Destroy empty container by perforation and never use for any other purpose.

**DIRECTIONS FOR USE:** Use only as directed.

Apply by hand, knapsak, or power duster.

| CROP                           | PEST                    | DOSAGE  | TIME OF APPLICATION                                |
|--------------------------------|-------------------------|---|--|
| <b>All crops</b>               | Astylus beetle          | 15-30 kg/ha   | Apply when pest appears and repeat when necessary. |
|                                | Lesser Armyworm         | 15-30 kg/ha   |  |
|                                | Armyworms               | 15 kg/ha  |  |
|                                | Chafer Beetles          | 15-25 kg/ha   |  |
|                                | Short-horn              | 15-30 kg/ha   |  |
|                                | Grasshopper             |   |  |
| <b>Tobacco</b>                 | Aphids, slug & Whitefly | Seedbed treatment<br>30 g/ 10m <sup>2</sup><br>Field treatment<br>15-25 kg/ha | Full cover application when necessary.             |
|                                |                         |   |  |
| <b>Sorghum</b>                 | Aphids                  | 15-30 kg/ha   | Full cover application when necessary.             |
| <b>Tomatoes</b>                |                         |   |  |
| <b>Flowers and Ornamentals</b> |                         |   |  |
| <b>Peaches and Plums</b>       | Aphids                  | 25 kg/ha  | Full cover application when necessary.             |
| <b>Groundnuts</b>              | Aphids                  | 10-15 kg/ha   | Full cover application when necessary              |
| <b>Maize</b>                   | Aphids                  | 12,5-15 kg/ha   |  |
| <b>Cotton</b>                  | Aphids                  | 30 kg/ha  | Full cover application when necessary.             |
| <b>Beans</b>                   | Aphids                  | 15-25 kg/ha   | Full cover application when pest is noticed.       |
|                                | CMR beetles             | 20-25 kg/ha   |  |
|                                | Thrips                  | 15-30 kg/ha   |  |

|  |   |             |  |
|--|---|-------------|--|
| <b>Flowers and</b>   | Thrips                                      | 15-30 kg/ha | Full application at weekly intervals as soon as Pest or damage is noticed. |
| <b>Ornamentals</b>   |   |             |  |
| <b>Onions</b>  | Thrips                                      | 20-30 kg/ha |  |
| <b>Lucerne</b>   | Caterpillar                                 | 10-30 kg/ha | Full cover application when pest is noticed.                               |
| <b>Mushrooms</b>   | Phorid and Sciarid Flies (Fungus gnats)     | 10-15 kg/ha | Apply thoroughly soon after picking. Repeat at weekly intervals.           |
| <b>Cruciferae (Cabbage, Cauliflower, Broccoli and Brussel sprouts)</b> | Greater Cabbage and Diamondback Moth Larvae | 20-30 kg/ha | Full cover application when pest is noticed. Repeat at weekly intervals.   |

**PUBLIC HEALTH**

| <b>SITE</b>                       | <b>PEST</b>       | <b>DOSAGE</b>                 | <b>APPLICATION DIRECTIONS</b>   |
|-----------------------------------|-------------------|-------------------------------|---|
| Garbage, Manure and Compost heaps | Fly-larvae        | Apply uniformly until visible | Full cover application to outside of heaps when fly breeding is noticed. Usually at weekly intervals. Apply again whenever fresh material is added. |
| Buildings, Rooms and Houses       | Fleas and bedbugs | Dust freely                   | Do not apply to the body. Apply freely to all places frequented by these pests.   |