

VOLCANO TRICLON

Reg. No.: L6661 Act No. 36 of 1947 • Namibia Reg. No.: N-AR 1605 • Botswana Reg. No.: W130241

A systemic emulsifiable concentrate herbicide for the control of woody plants and weeds as indicated for forestry, grass pastures and industrial areas.

HRAC HERBICIDE GROUP CODE: O

ACTIVE INGREDIENT:

Triclopyr (Pyridyloxy compound) 480 g/l
(as butoxyl ethyl ester)

DISTRIBUTOR:

PROTEK, a division of PE-BEE Agri (PTY) LTD
Co. Reg. No.: 2005/036308/07
P.O. Box 72, Heidelberg, 1438
Tel: (011) 812 9800 or 0861 PROTEK (0861 77 68 35)
www.proteksa.co.za

REGISTRATION HOLDER:

Arysta LifeScience South Africa (PTY) LTD
Co. Reg. No.: 2009/019713/07
7 Sunbury Office Park, Off Douglas Saunders Drive
La Lucia, 4019 • Tel: (031) 514-5600

EMERGENCY NO.: 082 446 8946

UN No.: 3082



HARMFUL

WARNINGS:

- Poisonous by swallowing.
- Concentrate may cause irritation to eyes and skin.
- Flammable.
- Store in a cool place.
- Store away from food, feed, seed and other agricultural chemicals and out of reach of children, uninformed persons and animals.
- May be toxic to fish.
- In case of poisoning, call a physician and consult first aid notes.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the weed to the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier or registration holder in the event of any uncertainty.

PRECAUTIONS:

For your own safety take the following precautions:

- Do not eat, drink or smoke while handling this product.
- Do not eat, drink or smoke before you have washed your hands, face and changed your clothing.
- After using this product, wash hands and face.
- Prevent contamination of food, feeds, drinking water and eating utensils.

When mixing:

- Wear eye protection and impermeable gloves.
- Wash with soap and running water after accidental skin contact.
- In case of eye contact, flush eyes immediately with clean running water. Seek medical advice.

When spraying:

- Do not inhale fumes or spray mist.
- Avoid contact with the spray solution.
- Wear suitable protective clothing.
- Prevent drift onto other crops, grazing, rivers, dams or areas not under treatment.

After spraying:

- Clean applicator thoroughly after use and dispose of rinsing water where it will not contaminate food, crops, grazing, rivers or dams.
- Change and wash your work clothes.
- Wash your exposed skin area and entire body, including your hair and under your fingernails, with soap and water.
- Rinse empty container three times with a volume of water equal to a minimum of 10 % of that of the container. Add the rinsing to the contents of the spray tank before destroying the container by perforation and flattening and NEVER use for any other purpose.

First aid:

Symptoms of poisoning:

Inhalation: Remove to fresh air if effects occur. Consult a physician.

Skin contact: Wash off in flowing water or shower.

Eye contact: Flush eyes with plenty of water.

Ingestion: Do not induce vomiting. Call a physician and/or transport to emergency facility immediately.

NOTE TO PHYSICIAN:

The decision of whether to induce vomiting or not should be made by an attending physician. If lavage is performed, endotracheal and/or esophageal control is suggested. Danger from lung aspiration must be weighed against toxicity when considering emptying the stomach. No specific antidote. Supportive care. Treatment based on judgement of the physician in response to reactions of the patient.

Use precautions:

Ensure that the person assigned to apply the product is properly trained in its use.

Resistance warning:

For resistance management **VOLCANO TRICLON** is a group code O herbicide. Any weed population may contain individuals naturally resistant to **VOLCANO TRICLON** and other group code O herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by **VOLCANO TRICLON** or any other group O herbicides.

To delay herbicide resistance:

1. Avoid exclusive repeated use of herbicides from the same herbicide group code. Alternate or tank mix with products from different herbicide codes.
2. Integrate control methods (Chemical, cultural and biological) into weed control programs. For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE: Use only as directed.

Compatibility:

VOLCANO TRICLON is compatible with ACTIPRON SUPER and GLYPHOSATE. In case of any uncertainty contact your nearest Volcano Triclon supplier.

Mixing:

For foliar applications, half fill the spray tank with water and add the required quantity of **VOLCANO TRICLON**. While filling the spray tank add ACTIPRON SUPER at the rate indicated below while maintaining constant agitation.

Application:

- Avoid drift.
- Do not spray under conditions when spray drift to desirable vegetation may occur.
- Always calibrate the equipment under field conditions.

A. Foliar application: (Use only on actively growing plants with full leaf cover)

- Apply as a full cover spray to leaves and stems.
- Do not apply to wet foliage.
- Avoid spray drift onto non-target plants, rivers, dams an untreated areas.
- Rain after application will have no effect on efficacy provided the spray has dried on the target plants.
- Use a knapsack sprayer with a solid cone nozzle (eg. Spraying Systems TG-1, Delavan CE 1 or equivalent type) that will ensure an even coverage of the target area. Maintain a constant pressure of between 200 – 250 kPa.

B. Cut stump application:

- Apply immediately or within 12 hours of felling.
- **VOLCANO TRICLON** in diesel should not be applied to stumps if surface is wet from rain or dew.
- Use solid cone nozzle tips (egg. Spraying systems TG-1, Delavan CE 1 or equivalent type) at 100 kPa (low pressure).
- Apply mixture to the point of run-off according to directions for stump and cut stumps in the table below.

C. Basal Bark Treatment: (for plants with stem diameter up to 10 cm)

- **VOLCANO TRICLON** in diesel should not be applied to stems if surface is wet from rain or dew.
- Apply by paintbrush or knapsack as a coarse low-pressure spray using narrow angle solid cone nozzle. Apply all round to the basal stem or trunk of the woody plant from soil level to a height indicated, as well as any exposed roots. In order to obtain satisfactory results a thorough wetting of the indicated area is imperative. Old or rough bark may require more mixture than young or smooth bark. In the case of multi-stemmed plants, each stem should be treated separately.

DOSAGE RATES AND METHOD OF APPLICATION:

A. FOLIAR APPLICATION:

NOTE: Actipron Super should be added at a rate of 50 ml/10 l spray mix for all foliar applications. Only apply to actively growing plants. Ensure thorough wetting to the point of runoff.

PLANT SPECIES	VOLCANO TRICLON / 10 l WATER	REMARKS
<i>Acacia cyclops</i> (Red Eye) <i>Acacia saligna</i> (Port Jackson willow)	50 ml (0,5 % mixture)	Only apply to young trees < 2 m in height.
<i>Acacia longifolia</i> (Long-leaved wattle)	60 ml (0,6 % mixture)	Only apply to young trees < 2 m in height.
<i>Acacia dealbata</i> (Silver wattle) <i>Acacia mearnsii</i> (Black Wattle)	25 – 75 ml (0,25 % – 0,75 % mixture)	Lower rate can be used on saplings < 50 cm. Higher rate on young trees 0,5 – 2 m tall.
<i>Asclepias fruticosa</i> (Shrubby milkweed)	50 ml (0,5 % mixture)	Ensure thorough wetting of all exposed plant parts.
<i>Caesalpinia decapetala</i> (Mauritius thorn)	50 ml (0,5 % mixture)	Apply to young trees up to maximum 3 m height. Taller plants must be cut back and regrowth treated only once it is 50 cm or higher
<i>Chromolaena odorata</i> (Triffid weed)	37,5 ml (0,375 % mixture)	Apply to young trees up to maximum 3 m height. Taller plants must be cut back and regrowth treated only once it is 50 cm or higher
<i>Eucalyptus grandis</i> (Bluegum)	75 ml (0,75 % mixture)	Apply to leaves and stems when coppice is between 1,0 and 1,5 m in height. Ensure that surfaces are thoroughly wetted. Repeat treatment if regrowth occurs.
<i>Rubus cuneifolius</i> (American bramble)	50 ml (0,5 % mixture)	Treat young actively growing trees, using 3 – 4 liter spray mixture per treated 100m ² . Large plants should be slashed and treated when regrowth reaches 0,5 m in height.
<i>Sesbania punicea</i> (Sesbania)	50 ml (0,5 % mixture)	Plants up to 2 m high may be treated. Taller plants must be cut back and regrowth treated when > 50 cm tall.
<i>Solanum mauritianum</i> (Bugweed)	50 ml (0,5 % mixture)	Plants < than 1,5 m (actively growing): Use 3 – 4 litre spraymix per treated 100 m ² . Plants > than 1,5 m: Slash and treat once re-growth is 0,5 m or higher.
<i>Solanum sisymbriifolium</i> (Wild tomato)	50 ml (0,5 % mixture)	Apply to actively growing plants until the point of runoff.
<i>Ulex europaeus</i> (Gorse)	100 ml (1,0 % mixture)	Treat young actively growing plants up to 1,0 m tall. Taller plants must be cut back and regrowth treated when at least 25 cm in height.

B. STUMP APPLICATION

PLANT SPECIES	VOLCANO TRICLON / 10 l DIESEL	REMARKS
<i>Acacia dealbata</i> (Silver wattle)	200 ml (2,0 % mixture)	Cut 10 – 15 cm above ground level. Apply to cut surface and all exposed bark and root crown of stumps.
<i>Cestrum laevigatum</i> (Inkberry)	100 ml (1,0 % mixture)	Cut stumps ± 10 cm above ground. Apply to cut surface and all exposed bark and roots.

PLANT SPECIES	VOLCANO TRICLON / 10 l DIESEL	REMARKS
<i>Chromolaena odorata</i> (Triffid weed)	100 ml (1,0 % mixture)	Cut stems ± 10 cm above ground. Apply to all cut surfaces and all exposed bark and roots.
<i>Eucalyptus grandis</i> <i>Eucalyptus diversicolor</i>	200 ml (2,0 % mixture)	Treat the cleaned, active growing region (cambium) of the cut surfaces, all the bark, root crown and any exposed roots. Spray to the point of runoff. Monitor the area and re-apply to coppice if necessary.
<i>Prosopis glandulosa</i> (Mesquite)	400 ml (4,0 % mixture)	Cut trees 10 – 15 cm above ground. Apply to cut stump (bark, root crown and exposed roots). Monitor the area and repeat steps above to all regrowth.
<i>Protasparagus laricinus</i> (Wild asparagus)	200 ml (2,0 % mixture)	Cut stems 10 – 15 cm above ground. Apply to cut stem surfaces and any exposed bark, root crown and roots. If any regrowth occurs, repeat steps above.
<i>Sesbania punicea</i> (Sesbania)	50 ml (0,5 % mixture) OR 100 ml in water (1,0 % mixture)	Cut stumps ± 10 cm above ground level. Apply to cut stem surfaces and any exposed bark, root crown and roots. For application in water only treat cut stem surfaces.

C. BASAL STEM APPLICATION

PLANT SPECIES	VOLCANO TRICLON / 10 l DIESEL	REMARKS
<i>Acacia decurrens</i> (Green wattle) <i>Acacia mearnsii</i> (Black wattle) <i>Acacia melanoxylon</i> (Blackwood)	200 ml (2,0 % mixture)	Ensure wetting of the root crown, exposed roots and stem up to a height of 0,25 m and ensure thorough wetting of all plant parts treated.
<i>Cestrum laevigatum</i> (Inkberry)	100 ml (1,0 % mixture)	Treat stems from ± 25 cm downward. Also treat exposed root crown and roots ensuring thorough wetting of all plant parts treated.
<i>Melia azedarach</i> (Syringa)	200 ml (2,0 % mixture)	Treat stems from ± 25 cm downward. Also treat exposed root crown and roots ensuring thorough wetting of all plant parts treated.
<i>Solanum mauritianum</i> (Bugweed)	200 ml (2,0 % mixture)	Treat stems from ± 25 cm downward. Also treat exposed root crown and roots ensuring thorough wetting of all plant parts treated.

VOLCANO TRICLON controls trees and shrubs as listed in the tables above. The registration holder does not accept any responsibility for unlisted trees and shrub species not being controlled or failure of the product due to application methods used, not recommended by the registration holder.

N.B. – In situations where more than one species of target plant occurs contact your local technical representative.

For assistance in the calibration for spot treatment please contact your local technical representative.

ACTIPRON SUPER (Mineral Oil) Reg. No. L 4923 is the Trademark of BP (SA) Ltd.

VOLCANO TRICLON

Reg. Nr.: L6661 Wet Nr. 36 van 1947 • Namibië Reg. Nr.: N-AR 1605 • Botswana Reg. Nr.: W130241

'n Sistemiese emulgeerbare konsentraat onkruidoder vir die beheer van houtagtige plante en onkruid soos aangetoon in bosbou, grasweidings en nywerheidsgebiede.

HRAC ONKRUIDODER GROEP KODE: O

AKTIEWE BESTANDEEL:

Triclopir (Piridiloksieverbinding) 480 g/l
(as butoksie-etielester)

VERSPREIDER:

PROTEK, 'n divisie van **PE-BEE Agri (EDMS) BPK**
Mpy. Reg. Nr.: 2005/036308/07
Posbus 72, Heidelberg, 1438
Tel: (011) 812 9800 of 0861 PROTEK (0861 77 68 35)
www.proteksa.co.za

REGISTRASIEHOUER:

Arysta LifeScience South Africa (EDMS) BPK
Mpy. Reg. Nr.: 2009/019713/07
7 Sunbury Office Park
Douglas Saunders-rylaan
La Lucia, 4019 • Tel: (031) 514-5600

NOOD NOMMER: 082 446 8946

UN Nr.: 3082



SKADELIK



WAARSKUWINGS:

- Giftig indien ingesluk word.
- Konsentraat mag irritasie van die oë en vel veroorsaak.
- Vlambaar.
- Stoor in 'n koel plek.
- Stoor weg van voedsal, voer, saad, ander landbou chemikalieë en buite bereik van kinders, oningeligte persone en diere.
- Mag giftig wees vir visse.
- In die geval van vergiftiging, raadpleeg 'n geneesheer en die eerstehulp nota's.

Ahoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie, aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale grond, klimaats en bergingstoestande, kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die onkruid teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer of registrasiehouer in die geval van enige onsekerheid.

VOORSORGMATREËLS:

Vir u eie veiligheid, maak gebruik van die volgende voorsorgmaatreëls:

- Moenie eet, drink of rook terwyl die produk hanteer word nie
- Moenie eet, drink of rook voordat u nie u hande en gesig gewas het nie, of voordat u nie skoon klere aangetrek het nie.
- Was hande en gesig na gebruik van die produk.
- Voorkom kontaminasie van kos, voere, drinkwater en eetgerei.

Tydens vermenging:

- Dra 'n beskermende bril en ondeurlaatsbare handskoene.
- In geval van toevallige kontak met vel, was onmiddellik met seep en lopende water.
- In geval van oogkontak, spoel oë met skoon lopende water uit en verkry onmiddellik mediese advies.

Tydens bespuiting:

- Moenie dampe of spuitnewel inasem nie.
- Vermoed kontak met die spuitmengsel.
- Dra geskikte beskermende klere.
- Verhoed spuitstofwegdrywing na ander gewasse, weidings, riviere, damme of gebiede wat nie behandel word nie.

Na bespuiting:

- Maak spuittoerusting wat gebruik is deeglik skoon en raak ontslae van spoelwater op so 'n plek wat dit nie voedsel, ander gewasse, weiding, riviere of damme sal besoedel nie.
- Wissel en was gekontamineerde klere.
- Was u hele liggaam, insluitende u hare en onder u vingernaels met seep en water.
- Spoel die houer driekeer uit met 'n volume water gelykstaande aan 'n minimum van 10 % van die inhoud van die houer. Gooi die spoelwater by die inhoud van die spuittenk voordat van die houer ontslae geraak word. Vernietig leë houer deur gate daarin te slaan en dit plat te slaan en moet vir geen ander doel hergebruik word nie.

Eerste Hulp maatreëls:

Simptome van vergiftiging:

Inaseming: Verwyder persoon na vars lug en raadpleeg 'n geneesheer.

Vel kontak: Was die besoedelde dele onder lopende water of stort.

Oog kontak: Spoel die oë MET GENOEGSAAM LOPENDE WATER.

Inname: Moet nie braking induuseer nie. Vervoer die persoon na die naaste geneesheer of nooddiens.

NOTA AAN DIE GENEESHEER:

Die besluitneming om braking te induser of nie, moet deur die waarnemende geneesheer gemaak word. Indien braking gedoen word, moet endotracheaal en of esophageale beheer toegepas word. Die gevaar van long aspirasie moet opgeweeg word teenoor die toksisiteit wanneer dit oorweeg word om die maag leeg te pomp.

Geen spesifieke teenmiddel bestaan nie. Ondersteunende behandeling moet toe gepas word. Behandeling moet gebaseer wees op die oordeel van die geneesheer in reaksie van die gedrag van die pasient.

Gebruiksvoorsorgmaatreël:

Verseker dat die toedieningsoperateur behoortlik in die toediening en hantering van die produk opgelei is. Bestandheidswaarskuwing **VOLCANO TRICLON** is 'n groepkode O onkruidoder. Enige populasie van 'n spesifieke onkruid mag individue insluit wat 'n natuurlike weerstand teen **VOLCANO TRICLON** of enige ander groepkode O onkruidodders het. Indien hierdie onkruidodders herhaalbaar aangewend word, kan die weerstandbiedende individue uiteindelik die onkruidpopulasie oorheers. Hierdie weerstandbiedende onkruid sal waarskynlik nie deur **VOLCANO TRICLON** of enige ander groepkode O onkruidoder beheer word nie.

Om weerstand teen onkruidodders te vertragen:

- Vermoed die eksklusiewe herhaalbare gebruik van onkruidodders met dieselfde groepkode.
- Wissel af met of gebruik tenkmengsels van produkte in verskillende onkruidoder groeppodes.
- Integreer ander beheermaatreëls (Chemies, verbouing en biologies) in onkruidoder programme.

Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

Verenigbaarheid:

VOLCANO TRICLON is verenigbaar met ACTIPRON SUPER en GLYPHOSATE. In die geval van enige onsekerheid, kontak u naaste **VOLCANO TRICLON** verskaffer.

Vermenging:

Vir looftoediening: maak die spuittenk half vol met water en voeg dan die verlangde volume **VOLCANO TRICLON** by. ACTIPRON SUPER moet bygevoeg word teen die aanbevole dosis soos hieronder aangedui terwyl die spuittenk gevul word. Maak seker dat die mengsel aanhoudend gerooer word.

Toediening:

- Verhoed spuitnewel wegdrywing.
- Moenie spuit in ongunstige toestande waar spuitnewel kan wegdrywe na nie-teiken plante.
- Maak seker dat die spuittoerusting korrek gekalibreer is onder veld toestande.

A. Loof toediening: (Wend slegs aan op aktief groeiende plante wat volle drag blare het.)

- Dien toe as 'n volle dekbespuiting op die blare en stamme.
- Moenie toedien terwyl die blare nat is nie.
- Verhoed spuitstofwegdrywing na gewasse, riviere, damme en gebiede wat nie behandel moet word nie.
- Moenie toedien as daar 'n moontlikheid van reën is nie. Reën na toediening sal geen invloed uitoefen op die effektiwiteit nie, indien daar genoeg tyd verloop het tussen toediening en reën sodat die spuitstof tyd gehad het om droog te word.
- Die spuitmengsel kan met 'n rugsakspuit toegedien word met soliede spuitpunt (Bv. Spraying Systems TG-1, Delavan CE1 of soortgelyke spuitstukke).
- Handhaaf 'n konstante druk van 200 – 250 kPa.

B. Gekapte stompbespuiting:

- Dien onmiddellik toe of verkieslik binne 12 uur na afkapping.
- **VOLCANO TRICLON** in diesel moet nie toegedien word op nat oppervlakte nie.
- Dien die spuitmengsel toe teen 'n lae druk (100 kPa) deur gebruik te maak van soliede keelspuitpunte (bv. Spraying Systems TG-1, Delavan of soortgelyke spuitstukke).
- Dien toe tot die punt van afdop volgens die aanwysings hieronder.

C. Stambespuiting: (vir plante met stamdeursnee tot 10 cm)

- **VOLCANO TRICLON** in diesel moet nie toegedien word op stamme indien hulle nat is weens reën of dou nie.
- Dien toe met 'n rugsakspuit met 'n smal soliede keelspuit as 'n growwe laedruk bespuiting. Dien toe rondom die stamme van houtagtige plante, vanaf grondhoogte tot 'n hoogte soos aangedui, asook op enige ontblote wortels. Deeglike benatting van die aangeduide areas is noodsaaklik vir goeie beheer. Ou of growwe bas mag meer spuitstof benodig as jonk of gladde bas. Waar meer as een stam per bos of struik voorkom, moet elke stam afsonderlik behandel word.

TOEDIENINGS HOEVEELHEDE EN METODE VAN TOEDIENING:

A. LOOF TOEDIENING

NOTA: Actipron Super moet teen 'n dosis van 50 ml/10 l spuitmengsel aangewend word vir alle looftoediening. Dien slegs toe op aktief groeiende plante. Verseker deeglike benatting deur tot die punt van afdop te benat.

PLANTSPESIE	VOLCANO TRICLON / 10 l WATER	OPMERKINGS
<i>Acacia cyclops</i> (Rooikrans) <i>Acacia saligna</i> (Port Jackson)	50 ml (0,5 % mengsel)	Behandel slegs jong bome < 2 m hoog.
<i>Acacia longifolia</i> (Langblaarwattel)	60 ml (0,6 % mengsel)	Behandel slegs jong bome < 2 m hoog.
<i>Acacia dealbata</i> (Silwerwattel) <i>Acacia mearnsii</i> (Swartwattel)	25 – 75 ml (0,25 % – 0,75 % mengsel)	'n Laer dosis kan gebruik word op saailinge < 50 cm. Hoër dosisse moet gebruik word op jong bome 0,5 – 2 m hoog.
<i>Asclepias fruticosa</i> (Tontelbossie)	50 ml (0,5 % mengsel)	Verseker deeglike benatting van alle plantdele.
<i>Caesalpinia decapetala</i> (Mauritiusdoring)	50 ml (0,5 % mengsel)	Behandel jong bome tot maksimum 3 m hoog. Hoër plante moet teruggesny word en hergroei moet behandel word op 50 cm of hoër.
<i>Chromolaena odorata</i> (Paraffienbos)	37,5 ml (0,375 % mengsel)	Behandel jong bome tot 'n maksimum van 3 m hoog. Hoër plante moet teruggesny word en hergroei moet behandel word op 50 cm of hoër.
<i>Eucalyptus grandis</i> (Bloekom)	75 ml (0,75 % mengsel)	Dien toe op blare en stamme sodra hergroei tussen 1,0 – 1,5 m hoog is. Maak seker dat die area deeglik benat word. Herhaal die toediening as hergroei weer voorkom.
<i>Rubus cuneifolius</i> (Amerikaanse braam)	50 ml (0,5 % mengsel)	Behandel slegs jong aktief groeiende boompies deur 3 – 4 liter spuitmengsel per bespuitte 100m ² te gebruik. Groter plante moet afgekap word en behandel word sodra die hergroei 0,5 m hoog is.
<i>Sesbania punicea</i> (sesbania)	50 ml (0,5 % mengsel)	Plante tot 2 meter hoog mag behandel word. Hoër plante moet teruggesny word en hergroei moet behandel word wanneer dit 50 cm hoog is.
<i>Solanum mauritianum</i> (Luisboom)	50 ml (0,5 % mengsel)	Plante < as 1,5 m: (aktief groeiend): Gebruik 3 – 4 liter spuitmengsel per bespuitte 100m ² . Plante > as 1,5 m: Kap af en behandel sodra die hergroei 0,5 m of hoër is.
<i>Solanum sisymbriifolium</i> (Wilde tamatie)	50 ml (0,5 % mengsel)	Behandel aktief groeiende plante deeglik tot by die punt van afdop.
<i>Ulex europaeus</i> (Brem)	100 ml (1,0 % mengsel)	Behandel aktief groeiende plante tot en met 1 m hoog. Groter plante moet eers teruggesny word. Sodra hergroei 25 cm in hoogte is kan plant behandel word.

B. STOMP TOEDIENING:

PLANTSPESIE	VOLCANO TRICLON / 10 l DIESEL	OPMERKINGS
<i>Acacia dealbata</i> (Silwerwattel)	200 ml (2,0 % mengsel)	Sny stompe af tot op 10 – 15 cm hoogte. Behandel die gesnyde oppervlakte asook alle bas, wortelkroon en blootgestelde wortels.
<i>Cestrum laevigatum</i> (Inkbessie)	100 ml (1,0 % mengsel)	Sny stompe af tot op 10 – 15 cm hoogte. Behandel die gesnyde oppervlakte asook alle bas, wortelkroon en blootgestelde wortels.

PLANTSPESIE	VOLCANO TRICLON / 10 l DIESEL	OPMERKINGS
<i>Chromolaena odorata</i> (Paraffienbos)	100 ml (1,0 % mengsel)	Sny stingels af tot op 10 – 15 cm hoogte. Behandel die gesnyde oppervlakte asook alle bas, wortelkroon en blootgestelde wortels.
<i>Eucalyptus grandis</i> <i>Eucalyptus diversicolor</i>	200 ml (2,0 % mengsel)	Behandel die skoon aktiewe (kambium) gedeelte van die gekapte oppervlakte sowel as die bas, wortelkroon en enige blootgestelde wortels. Spuit tot op die punt van afdop. Hou die bespuitte area dop en 'n opvolg behandeling mag nodig wees indien hergroei voorkom.
<i>Prosopis glandulosa</i> (Heuning prosopis)	400 ml (4,0 % mengsel)	Sny stompe af tot op 10 – 15 cm hoogte. Behandel die gesnyde oppervlakte asook alle bas, wortelkroon en blootgestelde wortels. Hou die bespuitte area dop en 'n opvolg behandeling soos bo mag nodig wees indien hergroei voorkom.
<i>Protasparagus laricinus</i> (Katbos)	200 ml (2,0 % mengsel)	Sny stompe af tot op 10 – 15 cm hoogte. Behandel die gesnyde oppervlakte asook alle bas, wortelkroon en blootgestelde wortels. Hou die bespuitte area dop en 'n opvolg behandeling soos bo mag nodig wees indien hergroei voorkom.
<i>Sesbania punicea</i> (Sesbania)	50 ml (0,5 % mengsel) OF 100 ml in water (1,0 % mengsel)	Sny stompe tot op 10 – 15 cm hoogte. Behandel gesnyde oppervlakte sowel as alle bas, wortelkroon en ontblote wortels. Vir behandelings in water behandel slegs gesnyde oppervlakte.

C. STAMBESPUITING

PLANTSPESIE	VOLCANO TRICLON / 10 l DIESEL	OPMERKINGS
<i>Acacia decurrens</i> (Groen wattel) <i>Acacia mearnsii</i> (Swart wattel) <i>Acacia melanoxylon</i> (Swarthout)	200 ml (2,0 % mengsel)	Behandel stamme tot op 'n hoogte van 25 cm. Verseker ook benatting van die wortelkroon en alle blootgestelde wortels.
<i>Cestrum laevigatum</i> (Inkbessie)	100 l (1,0 % mengsel)	Behandel stamme tot op 'n hoogte van 25 cm. Verseker ook benatting van die wortelkroon en alle blootgestelde wortels.
<i>Melia azedarach</i> (Sering)	200 l (2,0 % mengsel)	Behandel stamme tot op 'n hoogte van 25 cm. Verseker ook benatting van die wortelkroon en alle blootgestelde wortels.
<i>Solanum mauritianum</i> (Luisboom)	200 l (2,0 % mengsel)	Behandel stamme tot op 'n hoogte van 25 cm. Verseker ook benatting van die wortelkroon en alle blootgestelde wortels.

VOLCANO TRICLON beheer bome en struik volgens bogemelde etiket. Ander bome en struik wat nie tydens die ontwikkelingsproewe voorgekom het nie mag dalk ook beheer word of dan nie. Die registrasie houer aanvaar egter geen aanspreeklikheid teen opsigte van spesies wat nie op hierdie etiket gelys is nie, of toedienings metodes wat nie deur die registrasie houer aanbeveel is nie.

Belangrik: Wanneer meer as een ongewenste plantspesie voorkom raadpleeg u verskaffer of die registrasie houer.

Vir hulp met kalibrasie vir kolbehandeling raadpleeg u verskaffer of die registrasiehouer.

ACTIPRON SUPER (Mineral Oil) Reg. No. L 4923 is die Handelsmerk van BP (SA) BPK